

Quality improvement for community health services: A strategy to improve performance and quality in Kenya

There are over 6000 maternal deaths in Kenya every year - one of the highest rates in the world. Community health programmes have been shown to reduce maternal deaths and improve child survival. Community health workers (CHWs) undertake various tasks to support this, including case management of childhood illnesses (e.g. pneumonia, malaria, and neonatal sepsis), delivery of preventive interventions such as immunization, promotion of healthy behaviour, and mobilisation of communities. In Sub-Saharan Africa, CHW programmes have been an important strategy to address the severe shortages of skilled health workers, particularly for the poorest and most vulnerable populations.

Evidence has shown that appropriately trained CHWs have the potential to address barriers to preventive and curative care, to bring care closer to communities, and provide accessible services relevant to the needs at household level. However, if community-based services are to be effective, it is important that CHWs receive appropriate support that spans training in different aspects of community health and quality improvement methods as well as receiving regular support supervision and coaching beyond the initial training when they are recruited.

Kenya has had a national community strategy since 2006 with revisions in 2011 and 2014. The country set a target following the Recife workshop in Brazil in 2014 to increase the number of community health workers from 2,100 to 44,000 by 2017. Investment in expansion of community health programmes has however been limited with varying opinions by Kenyan policy makers and county leaders on their effectiveness. Rather than additional rapid expansion, a priority for the Ministry of Health in Kenya in 2017 is to improve the functionality of existing community units which requires that CHWs are appropriately trained, have the right tools, and report through national mechanisms.

With the devolution of health services in Kenya in 2013, counties became responsible for financing community health but this has varied widely with some counties considering CHWs important for achievement of health goals and investing in them while others focus on curative services. Decisions are made in the absence of data and evidence and are based on the priorities of local politicians. It is important to demonstrate effectiveness of community health and the significance of community health in the broader health

systems in order to gain local political support and more funding.

Globally, there has been interest in improving quality of health programmes with quality improvement (QI) gaining prominence. QI is a widely used management approach that engages multi-disciplinary teams from the bottom-up in developing solutions to resolve problems. QI interventions for community health, though few, have demonstrated results for maternal health. For example, in Tanzania more than 90% of women who had interacted with volunteers in a QI intervention village were going to a health facility for childbirth, compared with only around 60% of women in a non-QI intervention village.

In Kenya the updated Kenya Quality Model for Health (KQMH) was launched in 2014, QI teams have been active at facility level over the past six years with promising results. However until recently the KQMH model did not include community health services. In 2015 standards for community health services were published, however, robust mechanisms to track and support adherence to standards at community level have been lacking.



PHOTO BY Rosie Steege

www.reachoutconsortium.org

 twitter.com/REACHOUT_Tweet



This project is funded by the European Union.

Reachout
Linking Communities & Health Systems



Lessons learned

The key facilitators of QI among CHWs:

1. Supportive supervision of CHWs by their supervisors.

In sites where CHEWS and Sub-County supervisors offered regular supportive supervision sessions with CHWs as part of QI, there was a reported change in the quality of data collected by CHWs, an increase in the number of households covered by CHWs, and improved timeliness in reporting.

2. Training and continued mentorship and coaching on QI.

A one-off training was not enough to sustain learning. A phased approach with incremental learning proved effective. The participation of QI coaches during WIT meetings helped to reinforce the learning and correct implementation of QI approaches to improve quality of services.

3. Availability of tools and commodities.

Lack of commodities such as basic medicine, family planning commodities and HIV and malaria testing kits, by CHWs was reported to hinder delivery of good quality services and was a cause of frustration among community members. Lack of reporting tools directly affected the quality of reports interfering with the ability of CHWs and facilities to assess performance and improve services.

4. Team approach to implementation.

WITs had to be formed from among the CHWs, the CHEW, the Facility-in-Charge, and community members to strengthen referrals and community-facility linkage that were found to be a gap at the beginning of the intervention. Having a Facility-in-Charge as part of the community WIT was a motivator for CHWs who reported that it has improved the relationship between CHWs and the facility and served as a motivator for CHWs. The diversity of the team allowed different team members to take up responsibilities aligned to their strengths.

Recommendations for policy and practice

The study demonstrates that a QI approach at community level is both acceptable and feasible and can be implemented within the existing community health programmes. QI is not a new approach in Kenyan health system, however QI at community-level is novel. The project demonstrates that it enhances community linkage with the facility - a priority of community health programmes which has not always been successfully implemented.

QI approaches improve the quality of services delivered and reported by existing CHWs and their overall performance by improving their efficiency and effectiveness and contributes to overall achievement of County and Country primary health care goals.

Supervision and QI may not only improve performance of CHWs but also raise awareness of their role, legitimize their role in the community, and improve CHW motivation and retention.

1. Policy makers should ensure that QI is a priority in CHW policies, trainings, and funding priorities. This would enable the existing community health structures function optimally and achieve desired impact before having to invest in the more difficult process of increasing the number of CHWs.

2. By supporting implementation of QI for community health, health managers at County or district and facility level would benefit from improved results and motivated CHWs and community members who utilize services and are actively engaged. Managers should facilitate the process by providing the necessary tools, commodities, and ensuring supervision and feedback to and from CHWs and community members takes place consistently.

3. Alongside increased funding for community health services, donors need to prioritize QI in programme design to increase the efficiency, effectiveness, and impact of community health programmes.

Acknowledgements and funding

This research forms part of a multi-country research project, REACHOUT, a consortium aiming to maximize the equity, effectiveness and efficiency of close-to-community services in rural areas and urban slums in six countries (Bangladesh, Ethiopia, Indonesia, Kenya, Malawi, and Mozambique). The REACHOUT Consortium is funded by the European Union FP7 grant (number 306090). This document reflects only the author's views and the European Union is not liable for any use that may be made of the information contained therein.

The authors would like to acknowledge the assistance of the Community Health Development Unit in the Ministry of Health, Dr Salim, Daniel Kavoo and others. The research would not have been possible without the county and sub-county teams from Nairobi and Kitui, notably, Dr Ogara, Faith Kiruthi, Florence Opiyo, David Silu, Leonard Sharia Mbiyu, and Christine Sammy. Similarly we would like to thank the sub-county and community Work Improvement Teams, REACHOUT and USAID SQALE team members, and other LVCT Health staff.



Suggested citation:

Otiso L, Hawkins K, Muturi N, Mireku M, Doyle V, Taegtmeier M (2017) Quality improvement for community health services: A strategy to improve performance and quality in Kenya, REACHOUT Consortium Policy Brief

References

Haines A, Sanders D, Lehmann U, Rowe AK, Lawn JE, Jan S, Walker DG and Bhutta Z (2007) Achieving child survival goals: potential contribution of community health workers. *Lancet*. 2007 Jun 23;369(9579):2121-31

Horwood C, Butler L, Barker P, Phakathi S, Haskins L, Grant M and Rollins N (2017) A continuous quality improvement intervention to improve the effectiveness of community health workers providing care to mothers and children: a cluster randomised controlled trial in South Africa. *Human Resources for Health*, 15(1), 39. <https://doi.org/10.1186/s12960-017-0210-7>

Leatherman S, Ferris TG, Berwick D, Omaswa F, and Cris P (2010) The role of quality improvement in strengthening health systems in developing countries. *International Journal for Quality in Health Care*. <https://doi.org/10.1093/intqhc/mzq028>

McCullum R, Otiso L, Mireku M, Theobald S, De Koning K, Hussein S, and Taegtmeier M (2016) Exploring perceptions of community health policy in Kenya and identifying implications for policy change. *Health Policy and Planning*. <https://doi.org/10.1093/heapol/czv007>

Mireku M, Kiruki M, McCollum R, Taegtmeier M, de Koning K, and Otiso L (2014) Report on the Context Analysis of Close-to-Community Health Service Providers in Kenya.

Ministry of Health (2015) Kenya Quality Model for Health Quality Standards for Community Health Services. Level 1 2014. Retrieved from www.communityhealthservices.or.ke/images/pdf/2016_Documents/CHS-STANDARDS-final-version.pdf

Ministry of Health (2006) Taking the Kenya Essential Package for Health to the Community: A Strategy for the Delivery of Level One Services. Nairobi: Ministry of Health, pp. 1-60.

Petersen P B (1999) Total quality management and the Deming approach to quality management. *Journal of Management History*, 5, 468-488.

Tancred T, Manzi F, Schellenberg J, and Marchant T (2017) Facilitators and Barriers of Community-Level Quality Improvement for Maternal and Newborn Health in Tanzania. *Qualitative Health Research*. <https://doi.org/10.1177/1049732316638831>

CONTACT DETAILS

Lilian Otiso, LVCT Health, lilian.otiso@lvcthealth.org
www.lvcthealth.org